



18th European Weed Research Society Symposium

EWRS 2018

17-21 June 2018
Ljubljana, Slovenia

**New approaches for
smarter weed management**

www.ewrs2018.org

Book of Abstracts

Organiser

Kmetijski inštitut Slovenije (KIS) – Agricultural Institute of Slovenia, Hacquetova ulica 17, 1000 Ljubljana, Slovenija

Programme Committee

Dr. Jukka Salonen, Chairman / EWRS Vice-President; Prof. Dr. Hüsrev Mennan, Scientific Secretary; Prof. Dr. Paul Neve, EWRS President

EWRS Scientific Committee

Dr. Theo Verwijst, Department of Crop Production Ecology, Swedish University of Agricultural Sciences, Sweden; Prof. Dr. Garifalia Economou-Antonaka, Faculty of Crop Production Science, Agricultural University of Athens, Greece; Dr. Kirsten Torresen, Norwegian Institute of Bioeconomy Research (NIBIO), Norway; Dr. Roland Beffa, Bayer Crop Science AG, Integrated Weed Management & Resistance Biology, Germany; Dr. Ivo O. Brants, Monsanto Europe S.A, Belgium; Jan Petersen, University of Applied Science Bingen, Germany; Christian Bohren, Research Station Agroscope Changins-Wädenswil (ACW), Switzerland; Dr. Per Kudsk, Professor & Head of Section, Dept. of Agroecology, University of Aarhus, Denmark; Maurizio Vurro, Istituto di Scienze delle Produzioni Alimentari – CNR, Italy; Dr. Marleen Riemens, Wageningen University and Research Centre, Netherlands; Prof. Dr. Svend Christensen, University of Copenhagen, Faculty of Life Sciences, Denmark; Dr. Hanan Eizenberg, Newe Ya'ar Research Center, Israel; Dr. Euro Pannacci, Dept. of Agricultural, Food and Environmental Sciences – University of Perugia, Italy; Dr. Paula Westerman, Group Crop Health, Faculty of Agricultural and Environmental Science, University of Rostock, Germany

Local Organising Committee

Assoc. Prof. Dr. Andrej Simončič, President, Agricultural Institute of Slovenia; Dr. Robert Leskovšek, Agricultural Institute of Slovenia; Dr. Gregor Urek, Agricultural Institute of Slovenia; Prof. Dr. Stanislav Trdan, Biotechnical Faculty, University of Ljubljana; Prof. Dr. Mario Lešnik, Faculty of Agriculture and Life Sciences, University of Maribor; Ela Žilič, M. Sc., Agricultural Institute of Slovenia; Marjeta Urbančič Zemljič, M. Sc., Agricultural Institute of Slovenia

Editor

Andrej Simončič

Published by

Kmetijski inštitut Slovenije, 2018

The publication is published e-only – <http://www.ewrs.org>

The critical period of weed interference in upland rice in the Mid-West of Madagascar

Antsa Rafenomanjato¹, Antsa Rafenomanjato², Aude Ripoché³, Pascal Marnotte⁴, Paolo Barberi², Patrice Autfray³, Jean Augustin Randriamampianina⁵, Anna Camilla Moonen²

¹Scuola Superiore Sant' Anna, Pisa (Italy), ANTANANARIVO, Madagascar

²Institute of Life Sciences, Scuola Superiore Sant'Anna, PISA, Italy

³CIRAD, UPR AIDA, MONTPELLIER, France

⁴CIRAD UPR AIDA, SAINT-DENIS, Reunion

⁵FOFIFA, ANTANANARIVO, Madagascar

In lowland rice, weeds are generally well controlled by submergence, line transplanting and mechanical weeding. In upland rain-fed rice, weed control is highly challenging especially due to a peak workload at the beginning of the rainy season and weeding is done manually. This study aimed at determining the critical period of weed interference in a low-input upland rice cropping system, in order to optimize the timing of weeding. During the rainy season 2016-2017, a field experiment was conducted in Madagascar, Ivory station (19°33.29'S, 46° 24.913'E). Eight different weeding regimes were tested: a group of early weed interference (weedy until 20 DAS [days after sowing], 40 DAS and 60 DAS); a group of late weed interference (weedy from 20 DAS, 40 DAS and 60 DAS); two controls (weed-free and weedy). The experimental layout was a randomized complete block design with four replications. Data collected were weed cover at 20, 40 and 60 DAS, rice grain yield and weed biomass at harvest. The average rice yield in the weed-free control was 2.020 kg.ha⁻¹, and in the weedy control it dropped down to 10 kg.ha⁻¹ indicating more than 99% of yield loss. Early weed interference until 20, 40, and 60 DAS caused respectively yield losses of 6%, 60% and 87%; and late weed interference from 20, 40 and 60 DAS caused respectively yield losses of 63%, 39% and 9%. It indicated that weed presence before 20 DAS and after 60 DAS induced minor yield losses, while in-between these dates rice should be kept free from weed competition to avoid substantial losses. This information should advise farmers to better plan the timing of weeding, avoiding to invest labour in weeding when only a small increase in yield can be expected. This experiment will be repeated in the coming rainy season to confirm these results.